

## **An Official ERS Statement on Physical Activity in Chronic Obstructive Pulmonary Disease**

*Henrik Watz<sup>1\*</sup>, Fabio Pitta<sup>2</sup>, Carolyn L. Rochester<sup>3</sup>, Judith Garcia-Aymerich<sup>4</sup>, Richard ZuWallack<sup>5</sup>, Thierry Troosters<sup>6</sup>, Anouk W. Vaes<sup>7</sup>, Milo A. Puhon<sup>8</sup>, Melissa Jehn<sup>9</sup>, Michael I. Polkey<sup>10</sup>, Ioannis Vogiatzis<sup>11</sup>, Enrico M. Clini<sup>12</sup>, Michael Toth<sup>13</sup>, Elena Gimeno-Santos<sup>4</sup>, Benjamin Waschki<sup>1</sup>, Cristobal Esteban<sup>14</sup>, Maurice Hayot<sup>15/16</sup>, Richard Casaburi<sup>17</sup>, Janos Porszasz<sup>17</sup>, Edward McAuley<sup>18</sup>, Sally J. Singh<sup>19/20</sup>, Daniel Langer<sup>6</sup>, Emiel F.M. Wouters<sup>7/21</sup>, Helgo Magnussen<sup>1</sup>, and Martijn A. Spruit<sup>7/22\*</sup>*

*\*Task force co-chairs*

<sup>1</sup>Pulmonary Research Institute at LungenClinic Grosshandorf, Airway Research Center North, Member of the German Center for Lung Research, Grosshansdorf, Germany

<sup>2</sup>Laboratory of Research in Respiratory Physiotherapy, Department of Physiotherapy, Universidade Estadual de Londrina, Brazil

<sup>3</sup>Yale University School of Medicine, New Haven, CT, and VA Connecticut Healthcare System, USA

<sup>4</sup> Centre for Research in Environmental Epidemiology (CREAL); CIBER Epidemiología y Salud Pública (CIBERESP); and Universitat Pompeu Fabra, Departament de Ciències Experimentals i de la Salut, Barcelona, Spain

<sup>5</sup> St Francis Hospital Medical Center, Hartford, CT, USA

<sup>6</sup> Department of Rehabilitation Sciences, KU Leuven, Leuven, Belgium and Respiratory division and Pulmonary Rehabilitation, UZ Gasthuisberg, Leuven, Belgium

<sup>7</sup> Department of Research & Education, CIRO +, Center of expertise for chronic organ failure, Horn, the Netherlands

<sup>8</sup> Institute of Social and Preventive Medicine, University of Zurich, Switzerland

<sup>9</sup> Department of Pneumological Onkology and Transplantology, Charite Universitätsmedizin, Berlin, Germany

<sup>10</sup> NIHR Respiratory Biomedical Research Unit at the Royal Brompton & Harefield NHS Foundation Trust and Imperial College, London, UK

<sup>11</sup> National & Kapodistrian University of Athens, Medical School, 1<sup>st</sup> Department of Respiratory Medicine, Pulmonary Rehabilitation Unit, Athens, Greece

<sup>12</sup> Department of Medical and Surgical Sciences, University of Modena Reggio Emilia, Modena, Italy and Ospedale Villa Pineta, Pavullo n/F, Modena, Italy

<sup>13</sup> Department of Medicine, University of Vermont, College of Medicine, Burlington, VT, USA

<sup>14</sup> Pneumology Department, Hospital Galdakao-Usansolo, 48960 Galdakao, Bizkaia, Spain

<sup>15</sup> INSERM U-1046, University Montpellier I, Université Montpellier II, Montpellier, France

<sup>16</sup> CHRU Montpellier, Department of Clinical Physiology, F-34295 Montpellier, France

<sup>17</sup> Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, Los Angeles, California, USA

<sup>18</sup> Department of Kinesiology and Community Health, University of Illinois at Urbana-Champaign, Urbana, IL, USA

<sup>19</sup> Centre for exercise and rehabilitation Science, Glenfield Hospital, University Hospitals of Leicester NHS Trust, Leicester, UK

<sup>20</sup> Faculty of Health and Life Sciences , Coventry University, Coventry, UK

<sup>21</sup> Department of Respiratory Medicine, Maastricht University Medical Center (MUMC+), Maastricht, the Netherlands

<sup>22</sup> REVAL - Rehabilitation Research Center, BIOMED - Biomedical Research Institute, Faculty of Medicine and Life Sciences, Hasselt University, Diepenbeek, Belgium

Corresponding author:

Priv.-Doz. Dr. med Henrik Watz, MD

Pulmonary Research Institute at LungenClinic Grosshansdorf

Woehrendamm 80

D-22927 Grosshansdorf

Tel: +49-4102-8881-122

Fax: +49-4102-8881-113

Email: [h.watz@pulmoresearch.de](mailto:h.watz@pulmoresearch.de)

## **Abstract**

This European Respiratory Society (ERS) statement provides a comprehensive overview on physical activity in patients with chronic obstructive pulmonary disease (COPD).

Methods: A multidisciplinary Task Force of experts representing the ERS Scientific Group 01.02 'Rehabilitation & Chronic Care' determined the overall scope of this Statement through consensus. Focused literature reviews were conducted in key topic areas and the final content of this Statement was agreed upon by all members.

Results: The current knowledge regarding physical activity in COPD is presented, including the definition of physical activity, the consequences of physical inactivity on lung function decline and COPD incidence, physical activity assessment, prevalence of physical inactivity in COPD, clinical correlates of physical activity, effects of physical inactivity on hospitalizations and mortality, and treatment strategies to improve physical activity in patients with COPD.

Conclusions: This Task Force identified multiple major areas of research that need to be addressed further in the coming years, including, but not limited to, the disease-modifying potential of increased physical activity and to further understand how improvements in exercise capacity, dyspnea, and self-efficacy following interventions may translate into increased physical activity.

Recommendation: The Task Force recommends that this ERS Statement will be reviewed periodically (e.g. every 5-8 years).

## **1. Introduction**

Chronic obstructive pulmonary disease (COPD) is a highly prevalent chronic respiratory disease affecting about 10% of the adult population above 40 years of age (1). In addition to progressive chronic airflow limitation, patients with COPD commonly have multiple extra-pulmonary effects and comorbidities, which are associated with physical inactivity (2) (3). The “Global Initiative for Chronic Obstructive Lung Disease” has recommended regular physical activity for all COPD patients (2). However, the clinical relevance of regular physical activity has not been addressed in depth. The purpose of this Official European Respiratory Society (ERS) Statement is to highlight the existing science regarding physical (in)activity in patients with COPD, including, but not limited to, its prevalence, determinants, consequences, measurement, and potential treatment.

## **2. Methods**

An international group of pulmonologists, physiotherapists, movement scientists, exercise physiologists, health psychologists, social psychologists, and epidemiologists knowledgeable in the area of physical activity and/or COPD research was assembled into a Task Force by the leadership of the ERS Scientific Group 01.02 ‘Rehabilitation & Chronic Care’. Contributors searched the scientific literature (PubMed and Database of the Cochrane Collaboration) for original studies and systematic reviews relevant to the topic. Selection of relevant studies and reviews was based on the expertise of the contributors. Draft contributions were shared among committee members, reviewed and revised iteratively. Members were vetted for potential conflicts of interest according to the policies and procedures of ERS. This document represents the consensus of these Task Force members.

### 3. Definitions

Physical activity can be defined as any bodily movement produced by skeletal muscles that results in energy expenditure (4). Physical activity is a complex behavior that can be characterized by type, intensity, duration, patterns, and symptom experience. Exercise is a subset of physical activity. Exercise is physical activity that is planned, structured, repetitive, and purposeful (4). Physical activity also includes, but is not limited to, leisure-time, domestic, and occupational activities (4;5). Activities of daily living are another subset of physical activity and this term refers to a set of basic, everyday tasks required for personal self-care and independent living (6;7).

While physical inactivity can be defined simply as 'an absence of physical activity' (8), it is commonly used to represent a level of physical activity that is below an optimal or specified threshold. This concept rests on the strong evidence that lower levels of physical activity are related to poor health and predict negative health outcomes (9). In general, a healthy individual can be considered as physically inactive if either of the following criteria is not met: (1) 30 minutes of at least moderate-intensity physical activity on five or more days every week; (2) 20 minutes of vigorous-intensity physical activity on at least three days every week; or (3) an equivalent combination, which can also be accumulated in shorter bouts usually lasting 10 minutes of moderate (3 times 10 minutes) or vigorous (2 times 10 minutes) exercise (10;11). This represents a framework on which to base recommendations for physical activity promotion. However, the recommended intensity and duration of physical activity for the elderly may be different (12). Furthermore, the extent to which these recommendations apply to persons with COPD is currently unknown.

In recent years, attention has been raised regarding the adverse health effects of a sedentary lifestyle (13). A sedentary lifestyle is characterized by behaviors that do not increase skeletal muscle energy expenditure substantially above the resting level (14).

Sedentary individuals expend less than 10% of their total daily energy expenditure in performing moderate- or high-intensity activities (15).

#### **4. Consequences of physical inactivity in general population and in chronic diseases other than COPD**

Physical inactivity is a fundamental characteristic of many chronic diseases, both as a cause and as a consequence. Evidence suggests that reduced physical activity predisposes to greater incidence of cardiovascular disease (16;17), obesity (18), diabetes (19;20), cancer (21), dementia (22), and physical disability (23). In addition to its effect on development of chronic disease, physical inactivity may develop or worsen as a result of many diseases, owing to the sequelae of the disease and/or the associated reduction in physiological reserve. These effects of physical inactivity to both potentiate and develop from chronic disease likely explain the fact that reduced physical activity, measured either subjectively or objectively, is associated with higher overall mortality rates in the elderly (24;25). The exact mechanisms whereby physical inactivity interacts with anatomical and physiological changes related to aging and other pathological co-factors to contribute to the evolution of disease or subsequent morbidity or mortality, however, is poorly understood within the human population.

#### **5. Consequences of physical inactivity on lung function decline and COPD incidence**

Several population-based epidemiological studies have assessed the longitudinal effect of regular physical activity on lung function decline or COPD incidence (26-30); their main characteristics and results are summarised in Table 1. Briefly, all studies show an inverse relationship between physical activity levels and the magnitude of lung function decline in at least one of the population subgroups or physical activity variables studied. However, the association between lower physical activity levels and faster lung function decline is not

consistent across all population subgroups or physical activity variables. Potential explanations for such inconsistencies include selection bias, lack of adjustment for potential confounders, and lack of consideration of changes in physical activity level during follow-up. Only one study overcomes such limitations and, interestingly, shows beneficial effects of regular physical activity on lung function decline and COPD risk in active smokers but not in former or never smokers (29;30).

## **6. Physical activity assessment**

### *6.1 Questionnaires*

The physical activity of patients with COPD can be assessed using questionnaires. These instruments are commonly used in epidemiological studies and large clinical trials because they are inexpensive and easy to use (31). A variety of questionnaires exist that capture different aspects of physical activity such as amount, type, intensity, symptom experience, and limitations in the performance of 'activities of daily living' (31). The selection of a questionnaire to measure physical activity requires that the specific questionnaire fits the study aim, is properly developed and has strong psychometric properties, e.g., validation, test-retest reliability and responsiveness to change (32). In the specific case of COPD studies, additional criteria could be considered to improve validity of the questionnaire, such as the inclusion of information on low-intensity activities (31;33), or the availability of a version for interviewer-based administration (34). Other practical issues include the availability of a culturally validated version, the time required for questionnaire administration, and the ability to compare outcome levels across studies.

Recent systematic reviews have assessed all questionnaires available to measure physical activity in the elderly or chronically ill patients (35;36). From the 104 questionnaires identified in these systematic reviews, 15 were developed for use in patients with COPD. Validity was



assessed in 85% of these instruments, test-retest-reliability in 69%, responsiveness in only 19%, and none of the instruments was based on a conceptual framework (37). A current Innovative Medicines Initiative project is filling this gap by developing a valid Patient Reported Outcome tool (PROactive) capturing physical activity experience in COPD (<http://www.proactivecopd.com>).

A common methodological issue is recall bias, which may become a limitation if not addressed in the development and validation process. Garfield *et al* assessed 4 questionnaires against physical activity measured directly by accelerometry and found that while the Stanford Seven-Day Physical Activity Recall questionnaire could identify patients at both extremes of physical activity the other 3 had a poor relationship with directly measured activity only (38). Also in two other studies the relationship between questionnaire-derived physical activity measures and accelerometer-derived physical activity measures was either not given (39) or did not allow to reliably identify extremely inactive patients (40). .

Despite limitations of questionnaires when used on an individual level, some questionnaires might be used to measure physical activity in groups of patients with COPD. However, the choice depends on matching the question to be addressed with the psychometric properties of the instrument.

## 6.2 Step counters

Pedometers are small, lightweight, portable and nonintrusive devices which measure the number of steps performed in a given period of time. From that metric estimates of distance and energy expenditure can be made (31). Many pedometers are available, and variability exists not only in cost but also in mechanism of step detection, data storage, and sensitivity. Pedometers are most accurate at step counting, less accurate in distance estimates, and even less accurate at estimating energy expenditure (41). Pedometers may underestimate

the number of steps and energy expenditure during walking at slow speed, which is typical in patients with COPD (42-44). This may limit their accuracy in patients with moderate-to-very severe disease. However, pedometers may have positive role as a motivational tool aiming to increase daily activity, especially in addition to other combined interventions (45-47).

### *6.3 Activity monitors*

Accelerometers are electronic portable devices that are worn on the body to detect acceleration and thereby reflect bodily movement. They quantify activity counts, and may provide an estimate of time spent above or below a pre-specified activity level, number of steps, and energy expenditure (31). The use of accelerometers has received increasing interest since they add objective data which cannot be obtained from questionnaires or pedometers.

Accelerometers can detect movement along one axis (uni-axial accelerometers), two axes (bi-axial accelerometers), or three axes (tri-axial accelerometers). Uni-axial devices provide information similar to pedometers, but with the advantage of assessing acceleration in addition to simply detecting steps. Bi-axial and tri-axial devices allow for detection of movement in a wider range of physical activities, and are therefore more sensitive than uni-axial devices (48). Activity monitors sometimes combine accelerometers with other physiological sensors (such as heart rate, skin temperature) or are used in conjunction with positioning systems with the objective of increasing their accuracy to estimate daily physical activity and energy expenditure (49-51).

The validity of activity monitors for assessment of physical activity in patients with COPD has been the subject of many investigations in recent years (31;39;42;43;49;52-61). Two studies in COPD patients recently investigated the validity of six widely used accelerometers in comparison to the 'gold standards' of indirect calorimetry and doubly-labeled-water (62;63). Among the six devices, the DynaPort MiniMod , the Actigraph GT3X and the SenseWear Armband (all employing triaxial accelerometers) were valid and responsive for use in COPD

(62;63). These devices were demonstrated to be valid in other studies in COPD as well (42;43;49;52;54-56;58).

Several factors may influence the outcome of physical activity monitoring. Vibration from vehicle travel can falsely elevate activity counts measured by some devices, although this may be reduced by filtering the accelerometer signal (64). The number of assessment days and hours of use per day are also important factors that may influence the reliability of the physical activity assessment (65;66). Interestingly, compared to the other days of the week especially Sundays seem to be days of less physical activity in GOLD stage I to III patients (65). A recent study demonstrates that for cross-sectional analyses two to three days are sufficient for reliable measurement of physical activity in patients GOLD stage IV, whereas up to 5 days of measurement are required in patients with GOLD stage I (65). For measurements that aim to assess longitudinal changes 4 days were shown to be sufficient to demonstrate treatment effects following pulmonary rehabilitation in moderate to severe COPD, when weekends were excluded from the analysis (67).

The use of accelerometers certainly presents limitations. It should be noted that there is little uniformity in output from the various types of accelerometers, what makes it difficult to compare studies using different devices (68). Another limitation is that estimates of energy expenditure for individual patients may be inaccurate, especially among those with functional limitations due to chronic diseases that affect walking speed and efficiency of movement (69). Furthermore, purchase costs vary considerably between devices.

#### *6.4 Doubly labeled water*

The doubly labeled water (DLW) method provides an indirect assessment of total energy expenditure by the body over a substantial period of time (e.g. 2 weeks). The technique is described elsewhere (70). Briefly, known doses of deuterium ( $^2\text{H}_2\text{O}$ ) and  $^{18}\text{O}$  are ingested ( $^2\text{H}_2\text{O}$  &  $\text{H}_2^{18}\text{O}$ ). The deuterium washes out of the body through the urine, whereas the  $^{18}\text{O}$  is

eliminated as urine water and CO<sub>2</sub>. The difference between the wash out of the two (typically measured in urine samples) provides an estimate of CO<sub>2</sub> production by the body, which can be converted to energy expenditure (71).

The biggest drawback of the technique in the context of physical activity assessment in COPD is that it does not allow separating energy expenditure linked to physical activity and energy expenditure linked to basal metabolic rate or diet induced metabolism. Thus although the DLW technique is used to estimate total energy expenditure in patients with COPD(63;72;73) its ability to estimate physical activity is compromised by a number of assumptions that may be correct for healthy subjects but not for patients with COPD. For example one study used DLW in severe COPD and concluded that active energy expenditure was even higher in COPD compared to controls, which might be related to the increased oxygen cost of breathing and decreased mechanical efficiency in patients with COPD(72). On the other hand, actual physical activity levels are lower in patients with COPD (65). The DLW technique should likely be restricted to questions on caloric balance in patients with COPD rather than questions concerning the amount and intensity of physical activity. It can be questioned whether the technique should remain the 'gold standard' to validate monitors that measure the amount and intensity of movement in COPD, even though DLW will very likely remain a 'gold standard' for measuring the caloric cost of physical activity.

## **7. Levels of physical activity in patients with COPD**

Patients with COPD have significantly lower levels of physical activity as compared to healthy controls (65;74-81). Existing data shows that time spent walking is significantly lower in COPD patients compared to healthy, age-matched persons (74;76;77;80;82). These findings appear to be consistent across settings, cultural background, geographic area and methods used to measure physical activity. In addition, movement intensity of patients with

COPD is lower compared to age-matched healthy subjects, which indicates that patients with COPD walk at a slower pace (74). Current data suggest that patients with COPD reduce their physical activity early in the course of the disease (83-85). Accordingly, most patients do not meet currently recommended physical activity levels. For example in one study only 26% of 177 patients with a mean FEV<sub>1</sub> 52% predicted achieved at least 30 consecutive minutes of moderate intensity activity on at least 5 days; this increased to 50 % if the 30 minutes were accomplished in bouts of at least 10 minutes each (86). Corroborating these results, another study reported that only 29% of 73 patients with COPD achieved a mean of at least 30 minutes of moderate physical activity summed up throughout the day (87). Compared to controls COPD patients had a reduction by 50 % of their minutes of moderate physical activity per day, which was even more reduced when bouts of at least 10 min were compared between controls and COPD patients (88).

## **8. Factors associated with physical activity in COPD**

This section addresses associations between physical activity and clinical characteristics of patients with COPD, such as disease severity, comorbidities, exacerbations, and behavioral factors. Since most of the studies are cross-sectional it is not possible to draw conclusions regarding the directionality of the established associations (89). It should be acknowledged that in general, physical activity is dependent of many factors, which include biological, behavioral, genetic, social, environmental, cultural and policy factors (90). In this section we will discuss those aspects that are specifically studied in COPD.

### *8.1 Lung function*

FEV<sub>1</sub> shows a weak to moderate positive association with objectively measured physical activity in patients with COPD (39;65;74;77;79;80;91;92); Table 2). In general, FEV<sub>1</sub> explains only a small proportion of the variation in physical activity in subjects with COPD. Directly

measured maximal voluntary ventilation may provide an increased correlation with physical activity in this population (92). Fewer studies have investigated the relationship between physical activity and other lung function measures. While three studies found weak-to-moderate positive associations between physical activity and diffusion capacity (74;80;93), one study showed a strong and independent linear association (94). A robust inverse association was found between dynamic hyperinflation (measured in the laboratory during cardiopulmonary exercise test on a stationary cycle ergometer) and physical activity (95). Overall, while increasing severity of lung function impairment is associated with reduced physical activity in subjects with COPD, the relationship is relatively weak. Therefore, levels of physical activity cannot be accurately predicted from resting lung function parameters.

## *8.2 Exercise performance*

Physical fitness, which can be measured by various exercise tests, comprises a set of attributes that relates to the ability to perform physical activity (4). Accordingly, most studies in COPD have found moderately positive associations between either 6-minute walk distance or peak work rate in an incremental exercise test and objectively measured physical activity (40;65;74;79;96) (Table 3).

Two studies evaluated the predictive power of the 6-minute walk distance to identify physically inactive COPD patients with an objectively measured physical activity level lower than 1.4 (i.e. less than 40% of total daily energy expenditure is related to physical activity). In both studies 6-minute walk distance - even though moderately associated with physical activity - was found to be of limited value to reliably identify physically inactive COPD patients (40;65).

### *8.3 Self-efficacy*

Self-efficacy, an individual's belief in his or her capability of performing a specific task in a specific situation, is influenced by expectations of ability to perform the task and its outcome (97). Theoretically, higher levels of self-efficacy may be associated with increased physical activity; and higher levels of physical activity may result in enhanced self-efficacy belief. However, self-efficacy for walking, assessed in COPD using the Self-Efficacy Questionnaire – Walking (SEQ-W) instrument, was only weakly associated with objectively measured physical activity (91). Furthermore, in another study of 165 patients with COPD, general self-efficacy as measured with the General Self-Efficacy Scale (SES6) was not associated with physical activity (98).

### *8.4 Socio-demographic factors and environment*

Several socio-demographic and environmental factors including ethnicity, socioeconomic status, job availability, education level, patient choices about where to live, and seasonal variations in temperature and humidity have the potential to influence performance of daily physical activity among persons with COPD.

Among healthy adults, lower physical activity levels have been associated with lower socioeconomic status, lower education level and non-Caucasian race (99-101). However, this may not be the case for patients with COPD, since two studies have demonstrated a relationship between lower physical activity levels and higher socioeconomic status (102;103). In these studies, it is difficult to distinguish the impact of socioeconomic status from that of other influences, such as geographic location and ethnic and cultural differences, but it is possible that increased dependence on walking and public transportation among persons of lower socioeconomic status accounts for the findings. Also, both studies focused on persons with severe airflow limitation; hence the findings may not be applicable to patients with less severe lung function impairment.

Weather, climate and altitude conditions can also influence physical activity levels among persons with COPD. Extremes of heat or cold and or high environmental levels of particulates or other air pollutants can trigger increased symptoms, bronchoconstriction and acute exacerbations (104-106) and may pose a barrier to exercise and activity adherence (107;108). Accordingly, seasonal variations in daily physical activity have been reported (93;109-111), with a tendency for lower activity during periods with lower temperature. High altitude environments may also influence physical activity levels, since elevation in altitude leads to worsening resting and/or exercise hypoxemia and reduced exercise tolerance among patients with COPD (112). Geographical location *per se* may not strongly influence activity levels (e.g., independently of other factors such as climate, altitude or socioeconomic status) as no significant differences in physical activity were observed among patients with varying severity of COPD across diverse geographic locations in Europe and the United States (78;79). Finally, physical activity levels may be influenced by the day of the week; total activity and activity intensity were lower on Sundays or weekend as compared to other days (63;65).

### *8.5 Exacerbations of COPD*

Physical activity is dramatically reduced during and after hospitalization due to an exacerbation of COPD (113;114). Furthermore, recovery time is prolonged over several weeks and physical activity may not return to pre-exacerbation activity levels (113;114). Even patients with milder exacerbations, which do not require hospitalization, tend to stay indoors during the exacerbation period (115). Moreover, patients with a history of frequent exacerbations reduce their time spent outdoors at a faster rate compared with those with infrequent exacerbations, and thus are more likely to become housebound (115). This is confirmed by another study demonstrating that a history of more than one exacerbations is correlated with lower physical activity levels (79).

### *8.6 Comorbidities*



Comorbidities are common in COPD (116-118) and may independently impact physical activity levels. A cross-sectional study of 170 patients with COPD demonstrated that left ventricular cardiac dysfunction (assessed by elevated NT-pro-BNP and echocardiographic assessment of diastolic function) was associated with reduced physical activity levels, independent of COPD severity assessed by GOLD stage or the multidimensional BODE score (119). In this study, depression, anemia, systemic arterial blood pressure, and nutritional depletion were not associated with reduced physical activity levels (119). In the same cohort physical activity levels were significantly lower among patients with metabolic syndrome and COPD across all GOLD stages as compared with the patients without metabolic syndrome (120). It is not yet clear, which of the components of the metabolic syndrome contributed to this finding. For instance, obese COPD patients have lower activity levels compared to underweight- and normal weight patients (121;122). Furthermore, diabetes is strongly associated with inactivity in COPD, independent of other confounders (102). In a cohort of newly diagnosed COPD patients, physical inactivity was more strongly associated with the presence of comorbidities compared with airflow obstruction (123).

Quadriceps muscle strength and mass, which are commonly reduced in patients with COPD, are positively correlated with physical activity in several studies (74;79;85). In one study quadriceps muscle strength was predictive of physical activity, independent of FEV<sub>1</sub> (79). In contrast, handgrip strength did not correlate with physical activity in patients with COPD (94;119).

Mood disturbances such as symptoms of anxiety and depression are highly prevalent in patient with COPD (124). Of the six studies examining the relationship of physical activity and depression in patients with COPD (79;80;119;125-127) only two found significant associations (80;127). In one study higher levels of anxiety were associated with higher levels of physical activity whereas more depressive symptoms were associated with lower physical activity only when anxiety was part of the statistical model (127).

## 8.7 Systemic inflammation

In the general population a growing body of evidence demonstrates that physical activity, inflammation and immunity are tightly linked, with regular moderate physical activity reducing systemic inflammation (128-131). Potential mechanisms underlying these observations include the release of anti-inflammatory myokines by contracting skeletal muscles (132). In COPD, four studies demonstrated that higher levels of low-grade systemic inflammation are associated with lower levels of physical activity even after adjusting for relevant confounders (79;94;119;133). Whether anti-inflammatory mechanisms of regular physical activity or COPD-specific mechanisms of systemic inflammation underlie the associations of systemic inflammation and physical activity in COPD needs to be elucidated in further longitudinal studies (134).

## 8.8 Health status

Various assessments of health status have been related to physical activity measurements in patients with COPD. Most studies showed that an impaired health status assessed either by generic or disease-specific instruments is weakly to moderately related to a lower amount and intensity of physical activity (79;91;102;135;136). Changes in physical activity over time parallel trends in health status (137). Since most health status questionnaires have items addressing physical activity this could obviously affect these relationships.

## 8.9 Symptoms

Breathlessness on exertion is the primary symptom limiting exercise, which in turn leads to reduced physical activity in patients with COPD (138). Accordingly, patients' perception of breathlessness on exertion is reported to be a barrier for participation in exercise and daily activities of daily living (110;139). Indeed, greater levels of dyspnoea as measured by the modified Medical Research Council Dyspnoea Scale are related to lower levels of physical activity in patients with COPD (65;79).

Also fatigue is reported to be a frequent symptom in patients with COPD (140). There is one study that evaluated the association of fatigue assessed by the Functional Assessment of Chronic Illness Therapy - Fatigue questionnaire and physical activity. After adjusting for several confounders physical activity was demonstrated to be associated with fatigue in patients with COPD. (79). Whether and to what extent pain symptoms may affect physical activity in patients with COPD remains currently unknown.

### **9. Effect of physical inactivity on hospitalizations in patients with COPD**

To date seven prospective longitudinal studies have assessed the association between levels of regular physical activity and hospital admission or readmission due to COPD exacerbations (30;113;141-145). All studies consistently showed a statistically significant association of low physical activity levels with increased risk of hospitalizations. Although all studies reported associations adjusted for confounders, just one study adjusted for previous COPD hospitalizations, which is one of the most important risk factor for readmission (146). Importantly, the studies have shown that the amount of regular physical activity needed to obtain a significant effect on admissions due to COPD is relatively small, equivalent to walking or cycling 2 hours per week (30;113;141-145).

## **10. Effects of physical inactivity on survival in COPD**

Three longitudinal studies with a follow-up of 3 to 12 years have demonstrated that low levels of physical activity predict all-cause mortality in patients with COPD after controlling for relevant confounding factors (144;145;147). The relationship was consistent across differing settings, patient characteristics, and methods used to measure physical activity. In one study that separated respiratory-related from cardiovascular and all-cause mortality, the greatest effect was on the former (144). Recently, an assessment of physical activity has also been included as a prognostic factor in a multidimensional prognostic score for stable COPD patients (148).

Whether and to what extent improvements in physical activity levels may lower the risk of dying remains unknown in patients with COPD. Furthermore, it seems reasonable to hypothesize that patients who have a decline in physical activity over time have a worse prognosis compared to those patients who remain physically active.

## **11. Treatment strategies to improve physical activity**

### *11.1 Pharmacologic therapy*

While it is well-known that bronchodilators improve dyspnoea and exercise tolerance, only few studies have investigated the impact of bronchodilator therapy on physical activity (149-151). Two studies had positive results with regard to improvements of physical activity following bronchodilator treatment. Of these, one study (23 patients) was a non-randomized open-label study of a long-acting beta-agonist in a small number of patients (150), while the other study was a retrospective subgroup analysis (151). By contrast, two randomized, placebo-controlled, multicenter study could not demonstrate a change in physical activity following therapy with a long-acting bronchodilator (149;152). Therefore, it remains currently unknown, whether bronchodilator therapies that are known to improve exercise capacity will

also either improve physical activity or prevent deterioration of physical activity over time in patients with COPD.

### *11.2 Ambulatory oxygen therapy*

Oxygen therapy improves exercise tolerance in hypoxemic COPD patients, but whether activity levels are also enhanced is unclear. A small, randomized trial of replacing heavier oxygen tanks with lightweight ambulatory oxygen therapy failed to detect improvements in activity monitor assessments of physical activity over a 6 month period (153).

### *11.3 Pulmonary rehabilitation*

Pulmonary rehabilitation is a “comprehensive intervention to improve the physical and psychological condition of people with chronic respiratory disease and to promote the long-term adherence to health-enhancing behaviors” (154). Pulmonary rehabilitation has been demonstrated clearly to reduce dyspnea, increase exercise capacity and improve quality of life in patients with COPD. Increases in exercise capacity in combination with behavioral change may also have the potential to increase physical activity in patients with COPD. Despite this rationale the 10 studies that evaluated the effect of pulmonary rehabilitation on physical activity (80;82;155-162) have yielded inconsistent results; four showed an increase of physical activity (80;160-162) and six failed to do so (82;155-159). In the negative studies, the lack of improvement in physical activity was observed despite a concomitant increase in exercise capacity and quality of life. A systematic review and meta-analysis of single-group and randomized trials of the effect of exercise training (not necessarily pulmonary rehabilitation) on physical activity concluded that this intervention conferred a significant but small increase in this outcome (163).

### *11.4 Behavior changes, feedback, counseling*

Solely increasing the exercise capacity of patients with COPD may be insufficient to increase participation in self-directed leisure time activity (47;80;82;155;157-162;164). Widespread acknowledgment of the exercise maintenance problem has led to the identification of behavioral factors related to participation in daily physical activity and to the development of interventions targeting these factors.

Interventions including self-monitoring of activity behavior using activity monitors in combination with behavioral counseling in patients with COPD might have the potential to change physical activity behavior (45-47;165-167). However, the evidence base is weak since only few studies have been conducted using small samples (46;47) often without a control group (45;165;166). Larger studies with long term follow up are needed in patients with COPD.

Key components that increase the effectiveness of behavioral interventions have already been summarized in several meta-analyses and international guidelines (168-172) and include mobilizing social support, using well-described/established behavior change and self-regulatory techniques (self-monitoring, stimulus control, problem solving, relapse prevention management, goal setting, self-reinforcement, providing feedback on performance and developing action plans), providing higher contact time or contact frequency, and assessing the readiness / motivation to change (168;170). Motivational interviewing techniques have further been recommended as a collaborative communication approach (168).

#### 11.5 Summary of treatment strategies to improve physical activity

To date, only a few randomized controlled trials have studied the efficacy of pharmacologic or non-pharmacologic treatment strategies on daily physical activity in patients with COPD. Therefore, there is an urgent need for additional well-designed trials. Based on the correlates of physical activity, it seems reasonable to focus in future studies on physical, non-physical and environmental factors to improve physical activity in patients with COPD. Moreover,

future trials should not only focus on improvements of physical activity but also on the prevention of loss of physical activity.

## **12. Moving forward**

The scientific foundation regarding the clinical importance of assessing and improving physical activity in patients with COPD has grown considerably in the last decade.

Nevertheless, many questions remain unanswered and the methodology of physical activity assessment needs to be further standardized. Therefore, this Task Force identifies the following major areas that need to be further addressed in the coming years:

1. In addition to smoking cessation the disease-modifying potential of increased physical activity in smokers without COPD and in patients with all degrees of airflow limitation should be explored.
2. Further understanding is needed regarding the concepts to optimize the impact of the pharmacological and non-pharmacological interventions that aim to maintain or increase physical activity levels in patients with COPD.
3. Research should be undertaken to understand how improvements in exercise capacity, dyspnea, and self-efficacy following intervention (e.g. pulmonary rehabilitation, pharmacological therapy) might translate into increased physical activity.
4. The methodology to measure physical activity needs to be further standardized and formal guidance on that should be provided in future. Such methodologies could rely on objective and accurate assessment of physical activity, patient reported assessment of physical activity experience or combination thereof.

## **Acknowledgments**

The realization of the Official ERS Statement on Physical Activity in Chronic Obstructive Pulmonary Disease was not possible without the support of Prof. Dr. Guy Brusselle (ERS Guidelines Director), and Sandy Sutter (ERS CME & Guidelines Coordinator). Moreover, the Task Force co-chairs are grateful to the ERS for funding this ERS Statement.

Michael Polkey's contribution to this work was part funded by the NIHR Respiratory Biomedical Research Unit at the Royal Brompton & Harefield NHS Foundation Trust and Imperial College, who part fund his salary. Thierry Trooster's contribution was partly funded by the Flemish Research Foundation #G.0871.13. Anouk Vaes' contribution to this work was partially funded by 'Stichting de Weijerhorst' and Point-One funding from AgentschapNL, Dutch Ministry of Economic affairs, Netherlands. Martijn A. Spruit's contribution to this work was partially funded by Point-One funding from AgentschapNL, Dutch Ministry of Economic affairs, Netherlands. Benjamin Waschki's contribution to this work was partially funded by the German Center for Lung Research, Germany.



## Reference List

1. Buist AS, McBurnie MA, Vollmer WM, Gillespie S, Burney P, Mannino DM et al. International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet* 2007;370(9589):741-50.
  2. Vestbo J, Hurd SS, Agusti AG, Jones PW, Vogelmeier C, Anzueto A et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am.J.Respir.Crit Care Med.* 2013;187(4):347-65.
  3. Decramer M, Janssens W, Miravittles M. Chronic obstructive pulmonary disease. *Lancet* 2012;379(9823):1341-51.
  4. Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep.* 1985;100(2):126-31.
  5. Howley ET. Type of activity: resistance, aerobic and leisure versus occupational physical activity. *Med.Sci.Sports Exerc.* 2001;33(6 Suppl):S364-S369.
  6. Katz S. Assessing self-maintenance: activities of daily living, mobility, and instrumental activities of daily living. *J.Am.Geriatr.Soc.* 1983;31(12):721-7.
  7. Fricke J. Activities of Daily Living. In: JH Stone, M Blouin, editors. *International Encyclopedia of Rehabilitation*. Available online: <http://cirrie.buffalo.edu/encyclopedia/en/article/37/>. 2013.
  8. Organization W.H. Global Recommendation on Physical Activity for Health 2010. Available from: [http://apps.who.int/iris/bitstream/10665/44399/1/9789241599979\\_eng.pdf](http://apps.who.int/iris/bitstream/10665/44399/1/9789241599979_eng.pdf). 2013.
- Ref Type: Report
9. Wen CP, Wai JP, Tsai MK, Yang YC, Cheng TY, Lee MC et al. Minimum amount of physical activity for reduced mortality and extended life expectancy: a prospective cohort study. *Lancet* 2011;378(9798):1244-53.
  10. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA et al. Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association. *Circulation* 2007;116(9):1081-93.
  11. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* 2012;380(9838):247-57.
  12. Nelson ME, Rejeski WJ, Blair SN, Duncan PW, Judge JO, King AC et al. Physical activity and public health in older adults: recommendation from the American College of Sports Medicine and the American Heart Association. *Med.Sci.Sports Exerc.* 2007;39(8):1435-45.
  13. Owen N, Healy GN, Matthews CE, Dunstan DW. Too much sitting: the population health science of sedentary behavior. *Exerc.Sport Sci.Rev.* 2010;38(3):105-13.

14. Pate RR, O'Neill JR, Lobelo F. The evolving definition of "sedentary". *Exerc.Sport Sci.Rev.* 2008;36(4):173-8.
15. Bernstein MS, Morabia A, Sloutskis D. Definition and prevalence of sedentarism in an urban population. *Am.J.Public Health* 1999;89(6):862-7.
16. Li TY, Rana JS, Manson JE, Willett WC, Stampfer MJ, Colditz GA et al. Obesity as compared with physical activity in predicting risk of coronary heart disease in women. *Circulation* 2006;113(4):499-506.
17. Mora S, Cook N, Buring JE, Ridker PM, Lee IM. Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms. *Circulation* 2007;116(19):2110-8.
18. Li S, Zhao JH, Luan J, Ekelund U, Luben RN, Khaw KT et al. Physical activity attenuates the genetic predisposition to obesity in 20,000 men and women from EPIC-Norfolk prospective population study. *PLoS.Med.* 2010;7(8).
19. Sieverdes JC, Sui X, Lee DC, Church TS, McClain A, Hand GA et al. Physical activity, cardiorespiratory fitness and the incidence of type 2 diabetes in a prospective study of men. *Br.J.Sports Med.* 2010;44(4):238-44.
20. Jefferis BJ, Whincup PH, Lennon L, Wannamethee SG. Longitudinal associations between changes in physical activity and onset of type 2 diabetes in older British men: the influence of adiposity. *Diabetes Care* 2012;35(9):1876-83.
21. Friedenreich CM, Neilson HK, Lynch BM. State of the epidemiological evidence on physical activity and cancer prevention. *Eur.J.Cancer* 2010;46(14):2593-604.
22. Rovio S, Kareholt I, Helkala EL, Viitanen M, Winblad B, Tuomilehto J et al. Leisure-time physical activity at midlife and the risk of dementia and Alzheimer's disease. *Lancet Neurol.* 2005;4(11):705-11.
23. Gill TM, Allore HG, Gahbauer EA, Murphy TE. Change in disability after hospitalization or restricted activity in older persons. *JAMA* 2010;304(17):1919-28.
24. Manini TM, Everhart JE, Patel KV, Schoeller DA, Colbert LH, Visser M et al. Daily activity energy expenditure and mortality among older adults. *JAMA* 2006;296(2):171-9.
25. Matthews CE, George SM, Moore SC, Bowles HR, Blair A, Park Y et al. Amount of time spent in sedentary behaviors and cause-specific mortality in US adults. *Am.J.Clin.Nutr.* 2012;95(2):437-45.
26. Jakes RW, Day NE, Patel B, Khaw KT, Oakes S, Luben R et al. Physical inactivity is associated with lower forced expiratory volume in 1 second : European Prospective Investigation into Cancer-Norfolk Prospective Population Study. *Am.J.Epidemiol.* 2002;156(2):139-47.
27. Pelkonen M, Notkola IL, Lakka T, Tukiainen HO, Kivinen P, Nissinen A. Delaying decline in pulmonary function with physical activity: a 25-year follow-up. *Am.J.Respir.Crit Care Med.* 2003;168(4):494-9.

28. Cheng YJ, Macera CA, Addy CL, Sy FS, Wieland D, Blair SN. Effects of physical activity on exercise tests and respiratory function. *Br.J.Sports Med.* 2003;37(6):521-8.
29. Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity modifies smoking-related lung function decline and reduces risk of chronic obstructive pulmonary disease: a population-based cohort study. *Am.J.Respir.Crit Care Med.* 2007;175(5):458-63.
30. Garcia-Aymerich J, Lange P, Serra I, Schnohr P, Anto JM. Time-dependent confounding in the study of the effects of regular physical activity in chronic obstructive pulmonary disease: an application of the marginal structural model. *Ann.Epidemiol.* 2008;18(10):775-83.
31. Pitta F, Troosters T, Probst VS, Spruit MA, Decramer M, Gosselink R. Quantifying physical activity in daily life with questionnaires and motion sensors in COPD. *Eur.Respir.J.* 2006;27(5):1040-55.
32. Terwee CB, Mokkink LB, van Poppel MN, Chinapaw MJ, van Mechelen W, de Vet HC. Qualitative attributes and measurement properties of physical activity questionnaires: a checklist. *Sports Med.* 2010;40(7):525-37.
33. Forsen L, Loland NW, Vuillemin A, Chinapaw MJ, van Poppel MN, Mokkink LB et al. Self-administered physical activity questionnaires for the elderly: a systematic review of measurement properties. *Sports Med.* 2010;40(7):601-23.
34. Dinger MK, Oman RF, Taylor EL, Vesely SK, Able J. Stability and convergent validity of the Physical Activity Scale for the Elderly (PASE). *J.Sports Med.Phys.Fitness* 2004;44(2):186-92.
35. Frei A, Williams K, Vetsch A, Dobbels F, Jacobs L, Rudell K et al. A comprehensive systematic review of the development process of 104 patient-reported outcomes (PROs) for physical activity in chronically ill and elderly people. *Health Qual.Life Outcomes.* 2011;9:116.
36. Williams K, Frei A, Vetsch A, Dobbels F, Puhan MA, Rudell K. Patient-reported physical activity questionnaires: a systematic review of content and format. *Health Qual.Life Outcomes.* 2012;10:28.
37. Gimeno-Santos E, Frei A, Dobbels F, Rudell K, Puhan MA, Garcia-Aymerich J. Validity of instruments to measure physical activity may be questionable due to a lack of conceptual frameworks: a systematic review. *Health Qual.Life Outcomes.* 2011;9:86.
38. Garfield BE, Canavan JL, Smith CJ, Ingram KA, Fowler RP, Clark AL et al. Stanford Seven-Day Physical Activity Recall questionnaire in COPD. *Eur.Respir.J.* 2012;40(2):356-62.
39. Steele BG, Holt L, Belza B, Ferris S, Lakshminaryan S, Buchner DM. Quantitating physical activity in COPD using a triaxial accelerometer. *Chest* 2000;117(5):1359-67.
40. van Gestel AJ, Clarenbach CF, Stowhas AC, Rossi VA, Sievi NA, Camen G et al. Predicting daily physical activity in patients with chronic obstructive pulmonary disease. *PLoS.One.* 2012;7(11):e48081.
41. Schneider PL, Crouter SE, Lukajic O, Bassett DR, Jr. Accuracy and reliability of 10 pedometers for measuring steps over a 400-m walk. *Med.Sci.Sports Exerc.* 2003;35(10):1779-84.

42. Cavalheri V, Donaria L, Ferreira T, Finatti M, Camillo CA, Cipulo Ramos EM et al. Energy expenditure during daily activities as measured by two motion sensors in patients with COPD. *Respir.Med.* 2011;105(6):922-9.
43. Furlanetto KC, Bisca GW, Oldemberg N, Sant'anna TJ, Morakami FK, Camillo CA et al. Step counting and energy expenditure estimation in patients with chronic obstructive pulmonary disease and healthy elderly: accuracy of 2 motion sensors. *Arch.Phys.Med.Rehabil.* 2010;91(2):261-7.
44. Turner LJ, Houchen L, Williams J, Singh SJ. Reliability of pedometers to measure step counts in patients with chronic respiratory disease. *J.Cardiopulm.Rehabil.Prev.* 2012;32(5):284-91.
45. Moy ML, Weston NA, Wilson EJ, Hess ML, Richardson CR. A pilot study of an Internet walking program and pedometer in COPD. *Respir.Med.* 2012;106(9):1342-50.
46. Hospes G, Bossenbroek L, ten Hacken NH, van Hengel P, de Greef MH. Enhancement of daily physical activity increases physical fitness of outclinic COPD patients: results of an exercise counseling program. *Patient.Educ.Couns.* 2009;75(2):274-8.
47. de Blok BM, de Greef MH, ten Hacken NH, Sprenger SR, Postema K, Wempe JB. The effects of a lifestyle physical activity counseling program with feedback of a pedometer during pulmonary rehabilitation in patients with COPD: a pilot study. *Patient.Educ.Couns.* 2006;61(1):48-55.
48. Hikiyara Y, Tanaka S, Ohkawara K, Ishikawa-Takata K, Tabata I. Validation and comparison of 3 accelerometers for measuring physical activity intensity during nonlocomotive activities and locomotive movements. *J.Phys.Act.Health* 2012;9(7):935-43.
49. Patel SA, Benzo RP, Slivka WA, Sciurba FC. Activity monitoring and energy expenditure in COPD patients: a validation study. *COPD.* 2007;4(2):107-12.
50. Theou O, Jakobi JM, Vandervoort AA, Jones GR. A comparison of physical activity (PA) assessment tools across levels of frailty. *Arch.Gerontol.Geriatr.* 2012;54(3):e307-e314.
51. Troped PJ, Oliveira MS, Matthews CE, Cromley EK, Melly SJ, Craig BA. Prediction of activity mode with global positioning system and accelerometer data. *Med.Sci.Sports Exerc.* 2008;40(5):972-8.
52. Pitta F, Troosters T, Spruit MA, Decramer M, Gosselink R. Activity monitoring for assessment of physical activities in daily life in patients with chronic obstructive pulmonary disease. *Arch.Phys.Med.Rehabil.* 2005;86(10):1979-85.
53. Moy ML, Garshick E, Matthes KR, Lew R, Reilly JJ. Accuracy of uniaxial accelerometer in chronic obstructive pulmonary disease. *J.Rehabil.Res.Dev.* 2008;45(4):611-7.
54. Langer D, Gosselink R, Sena R, Burtin C, Decramer M, Troosters T. Validation of two activity monitors in patients with COPD. *Thorax* 2009;64(7):641-2.
55. Hill K, Dolmage TE, Woon L, Goldstein R, Brooks D. Measurement properties of the SenseWear armband in adults with chronic obstructive pulmonary disease. *Thorax* 2010;65(6):486-91.

56. Cohen MD, Cutaia M. A novel approach to measuring activity in chronic obstructive pulmonary disease: using 2 activity monitors to classify daily activity. *J.Cardiopulm.Rehabil.Prev.* 2010;30(3):186-94.
57. Sant'Anna T, Escobar VC, Fontana AD, Camillo CA, Hernandes NA, Pitta F. Evaluation of a new motion sensor in patients with chronic obstructive pulmonary disease. *Arch.Phys.Med.Rehabil.* 2012;93(12):2319-25.
58. Bauldoff GS, Ryan-Wenger NA, Diaz PT. Wrist actigraphy validation of exercise movement in COPD. *West J.Nurs.Res.* 2007;29(7):789-802.
59. Sugino A, Minakata Y, Kanda M, Akamatsu K, Koarai A, Hirano T et al. Validation of a compact motion sensor for the measurement of physical activity in patients with chronic obstructive pulmonary disease. *Respiration* 2012;83(4):300-7.
60. Annegarn J, Spruit MA, Uszko-Lencer NH, Vanbelle S, Savelberg HH, Schols AM et al. Objective physical activity assessment in patients with chronic organ failure: a validation study of a new single-unit activity monitor. *Arch.Phys.Med.Rehabil.* 2011;92(11):1852-7.
61. Van Remoortel H, Giavedoni S, Raste Y, Burtin C, Louvaris Z, Gimeno-Santos E et al. Validity of activity monitors in health and chronic disease: a systematic review. *Int.J.Behav.Nutr.Phys.Act.* 2012;9:84.
62. Van Remoortel H, Raste Y, Louvaris Z, Giavedoni S, Burtin C, Langer D et al. Validity of six activity monitors in chronic obstructive pulmonary disease: a comparison with indirect calorimetry. *PLoS.One.* 2012;7(6):e39198.
63. Rabinovich RA, Louvaris Z, Raste Y, Langer D, Remoortel HV, Giavedoni S et al. Validity of physical activity monitors during daily life in patients with COPD. *Eur.Respir.J.* 2013.
64. Cohen MD, Cutaia M, Brehm R, Brutus V, Pike VC, Lewendowski D. Detecting motor vehicle travel in accelerometer data. *COPD.* 2012;9(2):102-10.
65. Watz H, Waschki B, Meyer T, Magnussen H. Physical activity in patients with COPD. *Eur.Respir.J.* 2009;33(2):262-72.
66. Hecht A, Ma S, Porszasz J, Casaburi R. Methodology for using long-term accelerometry monitoring to describe daily activity patterns in COPD. *COPD.* 2009;6(2):121-9.
67. Demeyer H, Burtin C, Van Remoortel H, Hornikx M, Langer D, Decramer M et al. Standardizing the analysis of physical activity in patients with COPD following a pulmonary rehabilitation program. *Chest* 2014.
68. Butte NF, Ekelund U, Westerterp KR. Assessing physical activity using wearable monitors: measures of physical activity. *Med.Sci.Sports Exerc.* 2012;44(1 Suppl 1):S5-12.
69. Strath SJ, Pfeiffer KA, Whitt-Glover MC. Accelerometer use with children, older adults, and adults with functional limitations. *Med.Sci.Sports Exerc.* 2012;44(1 Suppl 1):S77-S85.

70. Westerterp KR, Wouters L, Marken Lichtenbelt WD. The Maastricht protocol for the measurement of body composition and energy expenditure with labeled water. *Obes.Res.* 1995;3 Suppl 1:49-57.
71. Schoeller DA. Recent advances from application of doubly labeled water to measurement of human energy expenditure. *J.Nutr.* 1999;129(10):1765-8.
72. Baarends EM, Schols AM, Pannemans DL, Westerterp KR, Wouters EF. Total free living energy expenditure in patients with severe chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 1997;155(2):549-54.
73. Arvidsson D, Slinde F, Nordenson A, Larsson S, Hulthen L. Validity of the ActiReg system in assessing energy requirement in chronic obstructive pulmonary disease patients. *Clin.Nutr.* 2006;25(1):68-74.
74. Pitta F, Troosters T, Spruit MA, Probst VS, Decramer M, Gosselink R. Characteristics of physical activities in daily life in chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 2005;171(9):972-7.
75. Schonhofer B, Ardes P, Geibel M, Kohler D, Jones PW. Evaluation of a movement detector to measure daily activity in patients with chronic lung disease. *Eur.Respir.J.* 1997;10(12):2814-9.
76. Singh S, Morgan MD. Activity monitors can detect brisk walking in patients with chronic obstructive pulmonary disease. *J.Cardiopulm.Rehabil.* 2001;21(3):143-8.
77. Hernandez NA, Teixeira DC, Probst VS, Brunetto AF, Ramos EM, Pitta F. Profile of the level of physical activity in the daily lives of patients with COPD in Brazil. *J.Bras.Pneumol.* 2009;35(10):949-56.
78. Troosters T, Sciurba F, Battaglia S, Langer D, Valluri SR, Martino L et al. Physical inactivity in patients with COPD, a controlled multi-center pilot-study. *Respir.Med.* 2010;104(7):1005-11.
79. Waschki B, Spruit MA, Watz H, Albert PS, Shrikrishna D, Groenen M et al. Physical activity monitoring in COPD: compliance and associations with clinical characteristics in a multicenter study. *Respir.Med.* 2012;106(4):522-30.
80. Walker PP, Burnett A, Flavahan PW, Calverley PM. Lower limb activity and its determinants in COPD. *Thorax* 2008;63(8):683-9.
81. Vorrink SN, Kort HS, Troosters T, Lammers JW. Level of daily physical activity in individuals with COPD compared with healthy controls. *Respir.Res.* 2011;12:33.
82. Coronado M, Janssens JP, de Muralt B, Terrier P, Schutz Y, Fitting JW. Walking activity measured by accelerometry during respiratory rehabilitation. *J.Cardiopulm.Rehabil.* 2003;23(5):357-64.
83. Van Remoortel H, Hornikx M, Demeyer H, Langer D, Burtin C, Decramer M et al. Daily physical activity in subjects with newly diagnosed COPD. *Thorax* 2013;68(10):962-3.

84. Gouzi F, Prefaut C, Abdellaoui A, Vuillemin A, Molinari N, Ninot G et al. Evidence of an early physical activity reduction in chronic obstructive pulmonary disease patients. *Arch.Phys.Med.Rehabil.* 2011;92(10):1611-7.
85. Shrikrishna D, Patel M, Tanner RJ, Seymour JM, Connolly BA, Puthuchearu ZA et al. Quadriceps wasting and physical inactivity in patients with COPD. *Eur.Respir.J.* 2012;40(5):1115-22.
86. Donaire-Gonzalez D, Gimeno-Santos E, Balcells E, Rodriguez DA, Farrero E, Batlle JD et al. Physical activity in COPD patients: patterns and bouts. *Eur.Respir.J.* 2012.
87. Vitorasso R, Camillo CA, Cavalheri V, Aparecida HN, Cortez VA, Sant'Anna T et al. Is walking in daily life a moderate intensity activity in patients with chronic obstructive pulmonary disease? *Eur.J.Phys.Rehabil.Med.* 2012;48(4):587-92.
88. Van Remoortel H, Camillo CA, Langer D, Hornikx M, Demeyer H, Burtin C et al. Moderate Intense Physical Activity Depends on Selected Metabolic Equivalent of Task (MET) Cut-Off and Type of Data Analysis. *PLoS.One.* 2013;8(12):e84365.
89. Gimeno-Santos E, Frei A, Steurer-Stey C, de Batlle J, Rabinovich RA, Raste Y et al. Determinants and outcomes of physical activity in patients with COPD: a systematic review. *Thorax* 2014.
90. Bauman AE, Reis RS, Sallis JF, Wells JC, Loos RJ, Martin BW. Correlates of physical activity: why are some people physically active and others not? *Lancet* 2012;380(9838):258-71.
91. Belza B, Steele BG, Hunziker J, Lakshminaryan S, Holt L, Buchner DM. Correlates of physical activity in chronic obstructive pulmonary disease. *Nurs.Res.* 2001;50(4):195-202.
92. Pitta F, Takaki MY, Oliveira NH, Sant'anna TJ, Fontana AD, Kovelis D et al. Relationship between pulmonary function and physical activity in daily life in patients with COPD. *Respir.Med.* 2008;102(8):1203-7.
93. Langer D, Iranzo MA, Burtin C, Verleden SE, Vanaudenaerde BM, Troosters T et al. Determinants of physical activity in daily life in candidates for lung transplantation. *Respir.Med.* 2012;106(5):747-54.
94. Garcia-Aymerich J, Serra I, Gomez FP, Farrero E, Balcells E, Rodriguez DA et al. Physical activity and clinical and functional status in COPD. *Chest* 2009;136(1):62-70.
95. Garcia-Rio F, Lores V, Mediano O, Rojo B, Hernanz A, Lopez-Collazo E et al. Daily physical activity in patients with chronic obstructive pulmonary disease is mainly associated with dynamic hyperinflation. *Am.J.Respir.Crit Care Med.* 2009;180(6):506-12.
96. Eliason G, Zakrisson AB, Piehl-Aulin K, Hurtig-Wennlof A. Physical activity patterns in patients in different stages of chronic obstructive pulmonary disease. *COPD.* 2011;8(5):369-74.
97. Strecher VJ, DeVellis BM, Becker MH, Rosenstock IM. The role of self-efficacy in achieving health behavior change. *Health Educ.Q.* 1986;13(1):73-92.

98. DePew ZS, Garofoli AC, Novotny PJ, Benzo RP. Screening for severe physical inactivity in chronic obstructive pulmonary disease: the value of simple measures and the validation of two physical activity questionnaires. *Chron.Respir.Dis.* 2013;10(1):19-27.
99. Marshall SJ, Jones DA, Ainsworth BE, Reis JP, Levy SS, Macera CA. Race/ethnicity, social class, and leisure-time physical inactivity. *Med.Sci.Sports Exerc.* 2007;39(1):44-51.
100. Crespo CJ, Ainsworth BE, Keteyian SJ, Heath GW, Smit E. Prevalence of physical inactivity and its relation to social class in U.S. adults: results from the Third National Health and Nutrition Examination Survey, 1988-1994. *Med.Sci.Sports Exerc.* 1999;31(12):1821-7.
101. Parks SE, Housemann RA, Brownson RC. Differential correlates of physical activity in urban and rural adults of various socioeconomic backgrounds in the United States. *J.Epidemiol.Community Health* 2003;57(1):29-35.
102. Garcia-Aymerich J, Felez MA, Escarabill J, Marrades RM, Morera J, Elosua R et al. Physical activity and its determinants in severe chronic obstructive pulmonary disease. *Med.Sci.Sports Exerc.* 2004;36(10):1667-73.
103. Pitta F, Breyer MK, Hernandez NA, Teixeira D, Sant'anna TJ, Fontana AD et al. Comparison of daily physical activity between COPD patients from Central Europe and South America. *Respir.Med.* 2009;103(3):421-6.
104. Koskela HO, Koskela AK, Tukiainen HO. Bronchoconstriction due to cold weather in COPD. The roles of direct airway effects and cutaneous reflex mechanisms. *Chest* 1996;110(3):632-6.
105. Wedzicha JA. Mechanisms of exacerbations. *Novartis.Found.Symp.* 2001;234:84-93.
106. Atkinson RW, Anderson HR, Sunyer J, Ayres J, Baccini M, Vonk JM et al. Acute effects of particulate air pollution on respiratory admissions: results from APHEA 2 project. *Air Pollution and Health: a European Approach.* *Am.J.Respir.Crit Care Med.* 2001;164(10 Pt 1):1860-6.
107. O'Shea SD, Taylor NF, Paratz JD. . . . But watch out for the weather: factors affecting adherence to progressive resistance exercise for persons with COPD. *J.Cardiopulm.Rehabil.Prev.* 2007;27(3):166-74.
108. Kosatsky T, Dufresne J, Richard L, Renouf A, Giannetti N, Bourbeau J et al. Heat awareness and response among Montreal residents with chronic cardiac and pulmonary disease. *Can.J.Public Health* 2009;100(3):237-40.
109. Moy ML, Danilack VA, Weston NA, Garshick E. Daily step counts in a US cohort with COPD. *Respir.Med.* 2012;106(7):962-9.
110. Katajisto M, Kupiainen H, Rantanen P, Lindqvist A, Kilpelainen M, Tikkanen H et al. Physical inactivity in COPD and increased patient perception of dyspnea. *Int.J.Chron.Obstruct.Pulmon.Dis.* 2012;7:743-55.
111. Sewell L, Singh SJ, Williams JE, Morgan MD. Seasonal variations affect physical activity and pulmonary rehabilitation outcomes. *J.Cardiopulm.Rehabil.Prev.* 2010;30(5):329-33.



112. Kelly PT, Swanney MP, Stanton JD, Frampton C, Peters MJ, Beckert LE. Resting and exercise response to altitude in patients with chronic obstructive pulmonary disease. *Aviat.Space Environ.Med.* 2009;80(2):102-7.
113. Pitta F, Troosters T, Probst VS, Spruit MA, Decramer M, Gosselink R. Physical activity and hospitalization for exacerbation of COPD. *Chest* 2006;129(3):536-44.
114. Borges RC, Carvalho CR. Physical activity in daily life in Brazilian COPD patients during and after exacerbation. *COPD.* 2012;9(6):596-602.
115. Donaldson GC, Wilkinson TM, Hurst JR, Perera WR, Wedzicha JA. Exacerbations and time spent outdoors in chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 2005;171(5):446-52.
116. Barnes PJ, Celli BR. Systemic manifestations and comorbidities of COPD. *Eur.Respir.J.* 2009;33(5):1165-85.
117. Nussbaumer-Ochsner Y, Rabe KF. Systemic manifestations of COPD. *Chest* 2011;139(1):165-73.
118. Vanfleteren LE, Spruit MA, Groenen M, Gaffron S, van Empel VP, Bruijnzeel PL et al. Clusters of comorbidities based on validated objective measurements and systemic inflammation in patients with chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 2013;187(7):728-35.
119. Watz H, Waschki B, Boehme C, Claussen M, Meyer T, Magnussen H. Extrapulmonary effects of chronic obstructive pulmonary disease on physical activity: a cross-sectional study. *Am.J.Respir.Crit Care Med.* 2008;177(7):743-51.
120. Watz H, Waschki B, Kirsten A, Muller KC, Kretschmar G, Meyer T et al. The metabolic syndrome in patients with chronic bronchitis and COPD: frequency and associated consequences for systemic inflammation and physical inactivity. *Chest* 2009;136(4):1039-46.
121. Monteiro F, Camillo CA, Vitorasso R, Sant'Anna T, Hernandez NA, Probst VS et al. Obesity and physical activity in the daily life of patients with COPD. *Lung* 2012;190(4):403-10.
122. Vozoris NT, O'Donnell DE. Prevalence, risk factors, activity limitation and health care utilization of an obese, population-based sample with chronic obstructive pulmonary disease. *Can.Respir.J.* 2012;19(3):e18-e24.
123. Van Remoortel H, Hornikx M, Langer D, Burtin C, Everaerts S, Verhamme P et al. Risk factors and comorbidities in the preclinical stages of chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 2014;189(1):30-8.
124. von Leupoldt A, Taube K, Lehmann K, Fritzsche A, Magnussen H. The impact of anxiety and depression on outcomes of pulmonary rehabilitation in patients with COPD. *Chest* 2011;140(3):730-6.
125. Moy ML, Matthess K, Stolzmann K, Reilly J, Garshick E. Free-living physical activity in COPD: assessment with accelerometer and activity checklist. *J.Rehabil.Res.Dev.* 2009;46(2):277-86.

126. Venkata A, DeDios A, ZuWallack R, Lahiri B. Are depressive symptoms related to physical inactivity in chronic obstructive pulmonary disease? *J.Cardiopulm.Rehabil.Prev.* 2012;32(6):405-9.
127. Nguyen HQ, Fan VS, Herting J, Lee J, Fu M, Chen Z et al. COPD Patients with Higher Levels of Anxiety Are More Physically Active. *Chest* 2013.
128. Handschin C, Spiegelman BM. The role of exercise and PGC1alpha in inflammation and chronic disease. *Nature* 2008;454(7203):463-9.
129. Febbraio MA. Exercise and inflammation. *J.Appl.Physiol* (1985.) 2007;103(1):376-7.
130. Gleeson M. Immune function in sport and exercise. *J.Appl.Physiol* (1985.) 2007;103(2):693-9.
131. Pedersen BK, Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat.Rev.Endocrinol.* 2012;8(8):457-65.
132. Pedersen BK. Exercise-induced myokines and their role in chronic diseases. *Brain Behav.Immun.* 2011;25(5):811-6.
133. Moy ML, Teylan M, Weston NA, Gagnon DR, Danilack VA, Garshick E. Daily Step Count is Associated with Plasma CRP and IL-6 in a US Cohort with COPD. *Chest* 2013.
134. Magnussen H, Watz H. Systemic inflammation in chronic obstructive pulmonary disease and asthma: relation with comorbidities. *Proc.Am.Thorac.Soc.* 2009;6(8):648-51.
135. McGlone S, Venn A, Walters EH, Wood-Baker R. Physical activity, spirometry and quality-of-life in chronic obstructive pulmonary disease. *COPD.* 2006;3(2):83-8.
136. Jehn M, Schindler C, Meyer A, Tamm M, Schmidt-Trucksass A, Stolz D. Daily walking intensity as a predictor of quality of life in patients with chronic obstructive pulmonary disease. *Med.Sci.Sports Exerc.* 2012;44(7):1212-8.
137. Esteban C, Quintana JM, Aburto M, Moraza J, Egurrola M, Perez-Izquierdo J et al. Impact of changes in physical activity on health-related quality of life among patients with COPD. *Eur.Respir.J.* 2010;36(2):292-300.
138. O'Donnell DE. Hyperinflation, dyspnea, and exercise intolerance in chronic obstructive pulmonary disease. *Proc.Am.Thorac.Soc.* 2006;3(2):180-4.
139. Bestall JC, Paul EA, Garrod R, Garnham R, Jones PW, Wedzicha JA. Usefulness of the Medical Research Council (MRC) dyspnoea scale as a measure of disability in patients with chronic obstructive pulmonary disease. *Thorax* 1999;54(7):581-6.
140. Todt K, Skargren E, Kentson M, Theander K, Jakobsson P, Unosson M. Experience of fatigue, and its relationship to physical capacity and disease severity in men and women with COPD. *Int.J.Chron.Obstruct.Pulmon.Dis.* 2014;9:17-25.
141. Benzo RP, Chang CC, Farrell MH, Kaplan R, Ries A, Martinez FJ et al. Physical activity, health status and risk of hospitalization in patients with severe chronic obstructive pulmonary disease. *Respiration* 2010;80(1):10-8.

142. Chen YJ, Narsavage GL. Factors related to chronic obstructive pulmonary disease readmission in Taiwan. *West J.Nurs.Res.* 2006;28(1):105-24.
143. Garcia-Aymerich J, Farrero E, Felez MA, Izquierdo J, Marrades RM, Anto JM. Risk factors of readmission to hospital for a COPD exacerbation: a prospective study. *Thorax* 2003;58(2):100-5.
144. Garcia-Aymerich J, Lange P, Benet M, Schnohr P, Anto JM. Regular physical activity reduces hospital admission and mortality in chronic obstructive pulmonary disease: a population based cohort study. *Thorax* 2006;61(9):772-8.
145. Garcia-Rio F, Rojo B, Casitas R, Lores V, Madero R, Romero D et al. Prognostic value of the objective measurement of daily physical activity in patients with COPD. *Chest* 2012;142(2):338-46.
146. Garcia-Aymerich J, Serra P, I, Mannino DM, Maas AK, Miller DP, Davis KJ. Lung function impairment, COPD hospitalisations and subsequent mortality. *Thorax* 2011;66(7):585-90.
147. Waschki B, Kirsten A, Holz O, Muller KC, Meyer T, Watz H et al. Physical activity is the strongest predictor of all-cause mortality in patients with COPD: a prospective cohort study. *Chest* 2011;140(2):331-42.
148. Esteban C, Quintana JM, Aburto M, Moraza J, Arostegui I, Espana PP et al. The health, activity, dyspnea, obstruction, age, and hospitalization: prognostic score for stable COPD patients. *Respir.Med.* 2011;105(11):1662-70.
149. O'Donnell DE, Casaburi R, Vincken W, Puente-Maestu L, Swales J, Lawrence D et al. Effect of indacaterol on exercise endurance and lung hyperinflation in COPD. *Respir.Med.* 2011;105(7):1030-6.
150. Hataji O, Naito M, Ito K, Watanabe F, Gabazza EC, Taguchi O. Indacaterol improves daily physical activity in patients with chronic obstructive pulmonary disease. *Int.J.Chron.Obstruct.Pulmon.Dis.* 2013;8:1-5.
151. Kesten S, Casaburi R, Kukafka D, Cooper CB. Improvement in self-reported exercise participation with the combination of tiotropium and rehabilitative exercise training in COPD patients. *Int.J.Chron.Obstruct.Pulmon.Dis.* 2008;3(1):127-36.
152. Troosters T, Sciurba FC, Decramer M, Siafakas NM, Klioze SS, Sutradhar SC et al. Tiotropium in patients with moderate COPD naive to maintenance therapy: a randomised placebo-controlled trial. *NPJ.Prim.Care Respir.Med.* 2014;24:14003.
153. Casaburi R, Porszasz J, Hecht A, Tiep B, Albert RK, Anthonisen NR et al. Influence of lightweight ambulatory oxygen on oxygen use and activity patterns of COPD patients receiving long-term oxygen therapy. *COPD.* 2012;9(1):3-11.
154. Spruit MA, Singh SJ, Garvey C, ZuWallack R, Nici L, Rochester C et al. An official american thoracic society/european respiratory society statement: key concepts and advances in pulmonary rehabilitation. *Am.J.Respir.Crit Care Med.* 2013;188(8):e13-e64.

155. Steele BG, Belza B, Hunziker J, Holt L, Legro M, Coppersmith J et al. Monitoring daily activity during pulmonary rehabilitation using a triaxial accelerometer. *J.Cardiopulm.Rehabil.* 2003;23(2):139-42.
156. Steele BG, Belza B, Cain KC, Coppersmith J, Lakshminarayan S, Howard J et al. A randomized clinical trial of an activity and exercise adherence intervention in chronic pulmonary disease. *Arch.Phys.Med.Rehabil.* 2008;89(3):404-12.
157. Dallas MI, McCusker C, Haggerty MC, Rochester CL, ZuWallack R. Using pedometers to monitor walking activity in outcome assessment for pulmonary rehabilitation. *Chron.Respir.Dis.* 2009;6(4):217-24.
158. Mador MJ, Patel AN, Nadler J. Effects of pulmonary rehabilitation on activity levels in patients with chronic obstructive pulmonary disease. *J.Cardiopulm.Rehabil.Prev.* 2011;31(1):52-9.
159. Egan C, Deering BM, Blake C, Fullen BM, McCormack NM, Spruit MA et al. Short term and long term effects of pulmonary rehabilitation on physical activity in COPD. *Respir.Med.* 2012;106(12):1671-9.
160. Sewell L, Singh SJ, Williams JE, Collier R, Morgan MD. Can individualized rehabilitation improve functional independence in elderly patients with COPD? *Chest* 2005;128(3):1194-200.
161. Mercken EM, Hageman GJ, Schols AM, Akkermans MA, Bast A, Wouters EF. Rehabilitation decreases exercise-induced oxidative stress in chronic obstructive pulmonary disease. *Am.J.Respir.Crit Care Med.* 2005;172(8):994-1001.
162. Pitta F, Troosters T, Probst VS, Langer D, Decramer M, Gosselink R. Are patients with COPD more active after pulmonary rehabilitation? *Chest* 2008;134(2):273-80.
163. Cindy Ng LW, Mackney J, Jenkins S, Hill K. Does exercise training change physical activity in people with COPD? A systematic review and meta-analysis. *Chron.Respir.Dis.* 2012;9(1):17-26.
164. Probst VS, Kovelis D, Hernandez NA, Camillo CA, Cavalheri V, Pitta F. Effects of 2 exercise training programs on physical activity in daily life in patients with COPD. *Respir.Care* 2011;56(11):1799-807.
165. Nguyen HQ, Gill DP, Wolpin S, Steele BG, Benditt JO. Pilot study of a cell phone-based exercise persistence intervention post-rehabilitation for COPD. *Int.J.Chron.Obstruct.Pulmon.Dis.* 2009;4:301-13.
166. Wewel AR, Gellermann I, Schwertfeger I, Morfeld M, Magnussen H, Jorres RA. Intervention by phone calls raises domiciliary activity and exercise capacity in patients with severe COPD. *Respir.Med.* 2008;102(1):20-6.
167. Vaes AW, Cheung A, Atakhorrami M, Groenen MT, Amft O, Franssen FM et al. Effect of 'activity monitor-based' counseling on physical activity and health-related outcomes in patients with chronic diseases: A systematic review and meta-analysis. *Ann.Med.* 2013;45(5-6):397-412.

168. Greaves CJ, Sheppard KE, Abraham C, Hardeman W, Roden M, Evans PH et al. Systematic review of reviews of intervention components associated with increased effectiveness in dietary and physical activity interventions. *BMC.Public Health* 2011;11:119.
169. Conn VS, Minor MA, Burks KJ, Rantz MJ, Pomeroy SH. Integrative review of physical activity intervention research with aging adults. *J.Am.Geriatr.Soc.* 2003;51(8):1159-68.
170. Conn VS, Hafdahl AR, Brown SA, Brown LM. Meta-analysis of patient education interventions to increase physical activity among chronically ill adults. *Patient.Educ.Couns.* 2008;70(2):157-72.
171. Estabrooks PA, Glasgow RE, Dzewaltowski DA. Physical activity promotion through primary care. *JAMA* 2003;289(22):2913-6.
172. Ferrier S, Blanchard CM, Vallis M, Giacomantonio N. Behavioural interventions to increase the physical activity of cardiac patients: a review. *Eur.J.Cardiovasc.Prev.Rehabil.* 2011;18(1):15-32.

Table 1. Studies on physical inactivity and lung function decline or COPD incidence.

Study	Design and setting	Number of subjects and follow-up	Subjects' characteristics	Physical activity assessment	Outcome(s)	Main Results
Jakes et al. [ref 26]	Population-based cohort  UK	n=12,283 mean 3.7 years follow-up	45% men  mean age: 59 (9) y  mean FEV <sub>1</sub> : 3.0 (0.7) L in men, and 2.2 (0.5) L in women	Standardised questionnaire	FEV <sub>1</sub> decline (mL/y)	<p>Climbing more stairs is associated with slower FEV<sub>1</sub> decline in women (-0.31, -0.11, -0.03, and 0.04% change in FEV<sub>1</sub> in none, 1-5, 6-10 and &gt;10 flights per day, p-for-trend&lt;0.004).</p> <p>Participation in vigorous leisure time activities is associated with slower FEV<sub>1</sub> decline in women (-0.22, -0.34, 0.27, and 0.03% change in FEV<sub>1</sub> in none, &lt;0.25, 0.25-1, and &gt;1 h per week, p-for-trend&lt;0.004).</p> <p>Based on a linear regression model adjusted for age, smoking status, and percent change in weight.</p> <p>No association found in men.</p>
Pelkonen et al. [ref 27]	Cohort of rural men with a high physical activity level	n=186  25 years follow-up	100% men  mean age: 54 (5) y  mean FEV <sub>0.75</sub> : 2.9 (0.6) L	Validated questionnaire	FEV <sub>0.75</sub> decline (mL/y)	<p>Decline in FEV<sub>0.75</sub> over 25 years was -44.4, -40.5, and -36.5 mL/y, p-for-trend=0.035 for low, middle, and high physical activity.</p> <p>Based on a linear regression model adjusted for age, height and</p>

	Finland					smoking habits.
Cheng et al. [ref 28]	Convenience sample from a clinic  USA	n=5707  mean 1.6 years follow- up	87% men  age range 25-55 years	Self-report	FEV <sub>1</sub> and FVC decline (mL/y)	Men who during the follow-up remained in the active category increased 50ml of FEV <sub>1</sub> and 70mL of FVC while subjects who remained in the sedentary group reduced 30 and 20 mL of FEV <sub>1</sub> and FVC, respectively.  Based on generalised least squared regressions adjusted for smoking and drinking habit change, age, baseline height, baseline lung function levels, and follow up time.  No association found in women.

Garcia-Aymerich et al. [ref 29]	Population-based cohort  Denmark	n=6790  10 years follow-up	43% men  mean age: 52 (12) y  mean FEV <sub>1</sub> : 2.7 (0.9) L  mean FVC: 3.3 (1.0) L	Validated questionnaire	COPD incidence (FEV <sub>1</sub> /FVC≤70 %)	<p>Active smokers with moderate to high physical activity show a reduced risk of developing COPD as compared to the low physical activity group (OR=0.77, p=0.027).</p> <p>Based on logistic regression models adjusted for gender, age, education, body mass index, weight change during follow-up, asthma, dyspnea, sputum, smoking status, and smoking duration.</p> <p>No association found in former or never smokers.</p>
---------------------------------	----------------------------------------	----------------------------------	------------------------------------------------------------------------------------------------------------	-------------------------	---------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------



					<p>FEV<sub>1</sub> and FVC decline (mL/y)</p> <p>Active smokers with moderate and high physical activity show a reduced FEV<sub>1</sub> decline compared to those with low physical activity (relative change of +2.6 and +4.8 mL/y of FEV<sub>1</sub>, p-for-trend=0.006)</p> <p>Active smokers with moderate and high physical activity show a reduced FVC decline compared to those with low physical activity (relative change of +2.6 and +7.7 mL/y of FVC, p-for-trend&lt;0.0001)</p> <p>Based on linear regression models adjusted for gender, age, education, body mass index, weight change during follow-up, ischemic heart disease, dyspnea, sputum, smoking status, smoking duration, alcohol consumption, and baseline lung function levels.</p> <p>No association found in former or never smokers.</p>
--	--	--	--	--	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Garcia-Aymerich et al. [ref 30]	Population-based cohort  Denmark	n=6568  mean 16 years follow-up	41% men  mean age: 49 (11) y  mean FEV <sub>1</sub> : 2.7 (0.8) L  mean FVC: 3.4 (1.0) L	Validated questionnaire	COPD incidence (FEV <sub>1</sub> /FVC≤70 %)	Subjects with moderate to high physical activity show a reduced risk of developing COPD as compared to the low physical activity group (OR=0.79, p=0.025).  Based on a logistic regression model adjusted for gender, age, education, body mass index, sputum, asthma and smoking, and weighted using marginal structural models to allow repeated measures of physical activity, lung function, and covariates.
					FEV <sub>1</sub> and FVC decline (mL/y)	Subjects with moderate and high physical activity show a reduced FEV <sub>1</sub> decline compared to those with low physical activity (relative change of +7.4 and +10.3 mL/y of FEV <sub>1</sub> , p-for-trend<0.001)  Subjects with moderate and high physical activity show a reduced FVC decline compared to those with low physical activity (relative change of +6.9 and +10.0 mL/y of FVC, p-for-trend<0.001)  Based on linear regression models adjusted for gender, age, education, body mass index, sputum, smoking, alcohol and baseline lung function levels, and weighted using marginal structural models to allow repeated measures of physical activity, lung function, and covariates.

Table 2. Relationship between FEV<sub>1</sub> and physical activity in some representative cross-sectional studies.

Study	Design and setting	Number of subjects (male/female)	Physical activity assessment	Main objective	R value	P value	Main results
Steele et al.  [ref 39]	Single-center  convenience sample  US	47 (44/3)	Accelerometer	Feasibility of accelerometer measurement	0.62	P<0.001	Preliminary data suggesting that a triaxial movement sensor is a reliable, valid, and stable measure of walking and daily physical activity in COPD patients.
Belza et al.  [ref 91 ]	Single-center  convenience sample  US	63(60/3)	Accelerometer	Evaluation of correlates of physical activity	0.37	P<0.01	Physical activity is most significantly related to walking abilities
Pitta et al.	Single-center	50 (36/14)	Accelerometer	Comparison of physical	0.28	p<0.05	Patients with COPD are markedly inactive in daily life.  Functional exercise capacity is the strongest correlate

[ref 74]	convenience sample Belgium			activity in COPD with healthy subjects			of physical activity.
Walker et al.  [ref 80]	Single- center  convenience sample  UK	23 (12/11)	Accelerometer	Evaluation of lower limb activity and the association of laboratory assessments with physical activity before and after rehabilitation in COPD	0.57	P<0.001	Physical activity in patients with COPD is closely related to leg activity, which is reduced compared with controls of similar age.
Hernandes  [ref 77]	Single- center  convenience	40 (18/22)	Accelerometer	Evaluation of physical activity and its	0.17	P=ns	Physical activity correlates only moderately with maximal and functional exercise capacity.

	sample  Brazil			clinical  correlates in  COPD in  Brazil			
Watz et al.  [ref 65]	Single-center  convenience sample  Germany	163 (122/41)	Accelerometer	Various  analyses of  physical activity and its clinical correlates in COPD	0.42	P<0.01	Clinical characteristics of patients with COPD only  incompletely reflect their physical activity
Waschki et al.  [ref 79]	Multi-center  convenience sample  UK, Netherlands	127(79/48)	Accelerometer	Evaluation of  compliance with wearing an accelerometer and the relationship of disease	0.65*	P<0.001	Excellent compliance with wearing a physical activity monitor in a multicenter study.  Consistent associations of physical activity with relevant disease characteristics in a multicenter study.

				characteristics with physical activity in a multi-center study			
--	--	--	--	----------------------------------------------------------------------------	--	--	--

\* Standardized regression coefficient beta adjusted for age, sex, study site, and BMI > 30

Table 3. Relationship between exercise tolerance and physical activity in some representative cross sectional studies.

Study	Design and setting	Number of subjects (male/female)	Main objective	Physical activity assessment	Exercise tolerance assessment	R value	P value	Main results
Steele et al. [ref 39]	Single-center convenience sample US	47 (44/3)	Feasibility of accelerometer measurement	Accelerometer	6MWD	0.74	P<0.001	Preliminary data suggesting that a triaxial movement sensor is a reliable, valid, and stable measure of walking and daily physical activity in COPD patients.

Belza et al.  [ref 91 ]	Single-center  convenience sample  US	63(60/3)	Evaluation of correlates of physical activity	Accelerometer	6MWD	0.60	P< 0.001	Physical activity is most significantly related to walking abilities
Pitta et al.  [ref 74]	Single-center  convenience sample  Belgium	50  (36/14)	Comparison of physical activity in COPD with healthy subjects	Acclerometer	6MWD, % pred.  Wmax, % pred.  Peak VO2, % pred.	0.76   0.64   0.33	P<0.0001   P<0.0001	Patients with COPD are markedly inactive in daily life.  Functional exercise capacity is the strongest correlate of physical activity.
Eliason et al.  [ref 96]	Multi-center  convenience sample  Sweden	44  (28/16)	Associations between physical activity and clinical characteristics of COPD	Accelerometer	6MWD	0.34*	P=0.03	Mean physical activity level and physical activity of at least moderate intensity are positively associated



								with exercise capacity
Waschki et al. [ref 79]	Multi-center convenience sample  UK, Netherlands	127  (79/48)	Evaluation of compliance with wearing an accelerometer and the relationship of disease characteristics with physical activity in a multi-center study	Accelerometer	6MWD	0.47**	P<0.001	Excellent compliance with wearing a physical activity monitor in a multicenter study.  Consistent associations of physical activity with relevant disease characteristics in a multicenter study.
Watz et al. [ref 65]	Single-center convenience sample  Germany	163  (122/41)	Various analyses of physical activity and its clinical correlates in COPD	Accelerometer	6MWD	0.46	P<0.001	Relation to clinical outcomes

Van Gestel et al. [ref 40]	Single-center convenience sample Switzerland	70 (49/21)	Predicting physical activity levels from 6MWD	Accelerometer	6MWD	0.69	P<0.001	6MWD cannot reliably predict physical inactivity
----------------------------	----------------------------------------------	------------	-----------------------------------------------	---------------	------	------	---------	--------------------------------------------------

6MWD, 6-minute walk distance; Wmax, maximal workload; peak VO<sub>2</sub>, peak oxygen uptake; \* standardized regression coefficient beta; \*\* standardized regression coefficient beta adjusted for age, sex, study site, and BMI > 30